FOR EXTERNAL ANIMAL USE ONLY

FIPROTEC® spot-on for dogs

Reg. No. G3794 Act 36/1947

CAUTION

INDICATIONS

For the treatment and prevention of flea and tick infestations on dogs and as an aid in the control of flea allergic dermatitis in dogs.

- Kills fleas and ticks following a single application.
- Residual action following a single dose:
  - Continues to kill fleas for up to 2 months (8 weeks).
  - Continues to kill ticks for up to 1 month (4 weeks).
- Monthly flea and tick treatment.

FIPROTEC® has a mode of action that effectively targets fleas and ticks. Fipronil collects in the oils of the skin and hair follicles and continues to be released onto the skin and coat, resulting in residual activity.

COMPOSITION

Fipronil 9.7 % m/v

STORAGE INSTRUCTIONS

Store in a cool dark place (below 25 ºC).

WARNINGS

- DO NOT TREAT PUPPIES YOUNGER THAN 10 WEEKS OF AGE.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Toxic to fish and bees.
- Avoid contact with hands or any other part of the body.
- Wash hands with soap and water after use.
- Do not smoke, eat or drink during application of the product.
- Dispose of used containers with domestic refuse.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- DO NOT bath or shampoo animals 3 days before or 3 days after application as this will decrease the duration of the activity.
- Remove a pipette from the rack.
- Twist off the top.
- Part the hair between the shoulder blades so that the skin is visible. Empty the contents of the container directly onto the visible skin by squeezing the pipette a number of times.
- If an excessive amount of the contents wet the actual hair it will impart a sticky appearance which will disappear within 24 hours.
- The active ingredient (Fipronil) will spread over the entire coat of the animal within 24 hours.
- Monthly application is recommended where there is a high risk of reinfestation, if the pet is allergic to fleas, or in situations where tick control is needed.

Application onto the skin

1 X pipette of 0.67 mℓ/dog up to 10 kg
1 X pipette of 1.34 mℓ/dog of 10 - 20 kg
1 X pipette of 2.68 mℓ/dog of 20 - 40 kg
1 X pipette of 4.02 mℓ/dog of 40 - 60 kg

Residual effect

A single dose will provide protection against fleas for up to 2 months and ticks for up to 1 month.

Registration holder:

Cipla Vet (Pty) Ltd
Co. Reg. No. 2001/017471/07
P O Box 1096, Durbanville, 7551
Tel: 0861 115 037

Cipla Vet
ANIMAL HEALTH

CD63A

SLEGS VIR UITWENDIGE DIEREGEBRUIK

FIPROTEC® spot-on vir honde

Reg. No. G3794 Wet 36/1947

VERSIGTIG

AANWYSINGS

Vir die behandeling en voorkoming van vlooi- en bosluisbesmetting op honde en as 'n hulpmiddel in die beheer van vlooi-allergiese dermatitis in honde.

- Dood vlooie en bosluise na 'n enkele behandeling.
- Nawerking na 'n enkele dosis:
  - Dood vlooie tot op 2 maande (8 weke).
  - Dood bosluise tot op 1 maand (4 weke).
- Maandelikse vlooi- en bosluisbehandeling.

FIPROTEC® het 'n meganisme van aksie wat vlooie en bosluise doeltreffend teiken. Fipronil versamel in die velolie en haarfollikels en hou aan om op die vel en haarkleed vrygelaat te word, met `n gevolglike nawerkende effek.

SAMESTELLING

Fipronil 9.7 % m/v

BERGINGSAANWYSINGS

Berg in 'n koel donker plek (benede 25 ºC).

WAARSKUWINGS

- MOENIE HONDJIES JONGER AS 10 WEKE OUD BEHANDEL NIE.
- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Giftig vir bye en visse.
- Vermy kontak met hande en enige ander gedeelte van die liggaam.
- Was hande met seep en water na gebruik.
- Moenie rook, eet of drink gedurende aanwending van die produk nie.
- Raak ontslae van houers saam met huishoudelike vullis.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- MOENIE honde 3 dae voor - of na behandeling bad of sjampoe nie; dit sal die tydsduur van aktiwiteit verminder.
- Maak 'n pipet los van die houer.
- Draai die prop af.
- Verdeel die hare tussen die skouerblaaie totdat die vel sigbaar is. Maak die inhoud van die houer leeg deur verskeie kere direk op die sigbare vel te druk.
- Voorsorg moet geneem word om die hare nie te nat te maak nie; dit sal die hare 'n klewering voorkoms gee by die plek van aanwending. Dit sal nietemin binne 24 ure na aanwending verdwyn.
- Die aktiewe bestanddeel (Fipronil) sal binne 24 ure oor die haarkleed versprei.
- Maandelikse toediening word aanbeveel waar daar 'n hoë risiko van herinfestasie is, as die dier allergies is vir vlooie, of waar bosluis beheer nodig is.

Aanwending op die vel

1 X pipet van 0.67 mℓ/hond tot 10 kg
1 X pipet van 1.34 mℓ/hond van 10 - 20 kg
1 X pipet van 2.68 mℓ/hond van 20 - 40 kg
1 X pipet van 4.02 mℓ/hond van 40 - 60 kg

Nawerking

'n Enkele dosis sal beskerming teen vlooie vir tot 2 maande en teen bosluise vir tot 1 maand verskaf.

Registrasiehouer:

Cipla Vet (Edms) Bpk
Mpy. Reg. Nr. 2001/017471/07
Posbus 1096, Durbanville, 7551
Tel: 0861 115 037

Cipla Vet
ANIMAL HEALTH

CD63A