

FOR EXTERNAL ANIMAL USE ONLY

FIPROTEC[®] spray

Reg. No. G3793 Act 36/1947

CAUTION

INDICATIONS

- For the treatment and prevention of flea and tick infestations on dogs and aids in the control of flea allergic dermatitis in dogs.
- Long acting flea and tick treatment for dogs.
- FIPROTEC[®] spray may be used on pregnant and lactating females and on unweaned puppies.

COMPOSITION

Fipronil 0.25g / 100ml

STORAGE INSTRUCTIONS

Store in a cool place.

WARNINGS

- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Toxic to bees and fish.
- Avoid contact with eyes.
- Wear gloves, especially during treatment of groups of animals.
- Spray in a well ventilated area.
- Wash hands immediately after use.
- Do not smoke, eat or drink during application.
- Keep your pet away from fires, heat or surfaces likely to be affected by the alcohol spray until dry.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- DO NOT bath or shampoo the animal 2 days before or 2 days after the application as this will decrease the duration of activity.
- Ruffle the dry coat against the lie of the hair before spraying.
- Spray the entire body of the animal at a distance of 10 - 20 cm.
- To ensure coverage of the head region spray FIPROTEC[®] spray onto a glove and rub gently around the face, avoiding the eyes.
- When your pet has been thoroughly covered, massage the coat all over to ensure that the spray gets right down to the skin.
- When treating puppies you may prefer to use a glove to apply FIPROTEC[®] spray.
- Allow your pet to dry naturally. Do not towel dry.
- Monthly applications are recommended where there is a high risk of re-infestation, if the pet is allergic to fleas and in situations where tick control is needed.

Application onto the skin

In order to wet the coat down to the skin it is recommended that the following application rates be used:

Short-haired animals (< 1.5 cm)

- A minimum of **3 ml/kg** body mass = 7,5 mg of active material per kg body mass.

Long-haired animals (> 1.5 cm)

- A minimum of **6 ml/kg** body mass = 15 mg of active material per kg body mass.

100 ml bottle

Each trigger application delivers 0.5 ml spray volume
e.g. for dogs under 12 kg

Short-haired animals (< 1.5 cm) apply **6** pump actions per kg (3 ml/kg)

Long-haired animals (> 1.5 cm) apply **12** pump actions per kg (6 ml/kg)

250 ml bottle

Each trigger application delivers 1 ml spray volume
e.g. for large dogs over 12 kg

Short-haired animals (< 1.5 cm) apply **3** pump actions per kg (3 ml/kg)

Long-haired animals (> 1.5 cm) apply **6** pump actions per kg (6 ml/kg)

Residual effect

Fleas: 2 months (1 - 3 months)

Ticks: 1 month (3 - 5 weeks)

This will depend on the parasite challenge in the animal's environment.

Registration holder:

Cipla Vet (Pty) Ltd
Co. Reg. No. 2001/017471/07
P O Box 1096, Durbanville, 7551
Tel: 0861 115 037



CD32 D

SLEGS VIR UITWENDIGE DIEREGBRUIK

FIPROTEC[®] sproei

Reg. No. G3793 Wet 36/1947

VERSIGTIG

AANWYSINGS

- Vir die behandeling en voorkoming van vlooi- en bosluisbesmetting op honde en as 'n hulpmiddel in die beheer van vlooi-allergie dermatitis by honde.
- Langwerkende vlooi- en bosluisbehandeling vir honde.
- FIPROTEC[®] sproei mag op dragtige- en lakterende diere sowel as ongespeende hondjies gebruik word.

SAMESTELLING

Fipronil 0.25g / 100ml

BERGINGSANWYSINGS

Berg in 'n koel plek.

WAARSKUWINGS

- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Giftig vir bye en visse.
- Vermoedlik met oë.
- Dra handskoene, veral gedurende behandeling van groepe diere.
- Gebruik slegs in 'n goed geventileerde area.
- Was hande dadelik na gebruik.
- Moenie rook, eet of drink gedurende gebruik nie.
- Hou die dier weg van vuur, hitte of oppervlaktes wat moontlik deur die alkohol-sproei geaffekteer kan word tot droog.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- MOENIE die dier 2 dae voor- of na behandeling bad of sjampoe nie, want dit sal die tydskedule van die aktiwiteit verminder.
- Roffel die droë haarkleed teen die lê van die hare voor aanwending.
- Sproei die hele lyf van die dier op 'n afstand van 10 - 20 cm.
- Om die kop te bedek, spuit FIPROTEC[®] sproei op 'n handskoen, vryf saggies oor die gesig en vermy die oë.
- Wanneer die dier deeglik benat is, vryf die haarkleed om te verseker dat die sproei tot op die vel penetreer.
- Wanneer hondjies behandel word, mag dit verkieslik wees om 'n handskoen te gebruik om FIPROTEC[®] sproei aan te wend.
- Laat die dier natuurlik droog word; moenie met 'n handdoek afdroog nie.
- Maandelikse toediening word aanbeveel waar daar 'n hoë risiko van herinfestasië is, as die dier allergies is vir vlooië, en waar bosluisbeheer nodig is.

Aanwending op die vel:

Om die haarkleed tot teen die vel nat te maak word dit aanbeveel dat aanwending teen die volgende skaal gedoen word:

Korthaar diere (< 1.5 cm)

- 'n Minimum van **3 ml/kg** liggaamsmassa = 7.5 mg aktiewe bestanddeel per kg liggaamsmassa.

Langhaar diere (> 1.5 cm)

- 'n Minimum van **6 ml/kg** liggaamsmassa = 15 mg aktiewe bestanddeel per kg liggaamsmassa.

100 ml bottel

Elke snellertrekking lewer 0.5 ml sproei volume
bv. vir honde onder 12 kg

Korthaar diere (< 1.5 cm) wend **6** pompaksies per kg (3 ml/kg) aan

Langhaar diere (> 1.5 cm) wend **12** pompaksies per kg (6 ml/kg) aan

250 ml bottel

Elke snellertrekking lewer 1 ml sproei volume
bv. vir groot honde meer as 12 kg

Korthaar diere (< 1.5 cm) wend **3** pompaksies per kg (3 ml/kg) aan

Langhaar diere (> 1.5 cm) wend **6** pompaksies per kg (6 ml/kg) aan

Nawerking

Vlooië: 2 maande (1 - 3 maande)

Bosluise: 1 maand (3 - 5 weke)

Dit sal afhang van die parasietuitdaging in die dier se omgewing.

Registrasiehouer:

Cipla Vet (Edms) Bpk
Mpy. Reg. Nr. 2001/017471/07
Posbus 1096, Durbanville, 7551
Tel: 0861 115 037



CD32 D