

FOR ANIMAL USE ONLY

SLEGS VIR DIEREGERBRUIK

TRIWORM-D MEDIUM DOG INDIVIDUAL

Reg. No. G3645 Act/Wet 36/1947

Worm remedy against ascarids, hookworms, whipworms and tapeworms in dogs
Wurmmiddel teen askariedes, haakwurms, sambokwurms en lintwurms in honde

STORAGE INSTRUCTIONS/BERGINGSAAWYSINGS

Store below/Berg onder 25 °C.

EACH TABLET CONTAINS/ELKE TABLET BEVAT:

Praziquantel/Prasikwantel 50 mg
Pyrantel pamoate/Pirantelpamoat 144 mg
Febantel 150 mg

CONTENT / INHOUD

2 Tablets / 2 Tablette

FOR FULL INSTRUCTIONS SEE ENCLOSED PACKAGE INSERT
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUBILJET

Registration holder/Registrasiehouer:

Ascendis Vet (Pty) Ltd
Co. Reg. No. 2001/017471/07
31 Georgian Crescent East
Bryanston
Johannesburg
South Africa
2191
Tel: [+27 10 880 0778](tel:+27108800778)
www.ascendisanimalhealth.com

Lot. No.:
Expires/Verval:

BAR CODE

PACKAGE INSERT

<p>FOR ANIMAL USE ONLY</p> <p>TRIWORM-D MEDIUM DOG INDIVIDUAL Reg. No. G3645 Act 36/1947</p> <p>INDICATIONS Worm remedy against ascarids, hookworms, whipworms and tapeworms in dogs.</p> <p>COMPOSITION Each tablet contains: Praziquantel 50 mg Pyrantel pamoate 144 mg Febantel 150 mg</p> <p>STORAGE INSTRUCTIONS Store below 25 °C.</p> <p>WARNINGS</p> <ul style="list-style-type: none">• DO NOT USE SIMULTANEOUSLY WITH PIPERAZINE COMPOUNDS.• Keep out of reach of uninformed persons, children and animals.• Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder. <p>DIRECTIONS FOR USE: USE ONLY AS DIRECTED</p> <ul style="list-style-type: none">• Can be administered directly, or mixed with meat, or crushed and mixed with feed.• Feeding and fasting before treatment is not necessary.	<p>SLEGS VIR DIEREGERBRUIK</p> <p>TRIWORM-D MEDIUM HOND ENKEL EENHEID Reg. No. G3645 Wet 36/1947</p> <p>AANWYSINGS Wurmmiddel teen askariedes, haakwurms, sambokwurms en lintwurms in honde</p> <p>SAMESTELLING Elke tablet bevat: Prasikwantel 50 mg Pirantelpamoaat 144 mg Febantel 150 mg</p> <p>BERGINGSAAWYSINGS Berg onder 25 °C.</p> <p>WAARSKUWINGS</p> <ul style="list-style-type: none">• MOENIE SAAM MET PIPERASIEN SAMESTELLINGS GEBRUIK NIE.• Hou buite bereik van oningeligte persone, kinders en diere.• Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer. <p>GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI</p> <ul style="list-style-type: none">• Kan direk toegedien word, óf met vleis gemeng, óf verkrummel en met kos gemeng.• Dit is nie nodig om diere te voer of uit te honger voor toediening nie.
<p>Dosage Dose according to body mass i.e. 1 tablet/10 kg e.g.</p>	<p>Dosis Doseer volgens liggaamsmassa d.w.s. 1 tablet/10 kg b.v.</p>

Size	Body mass/ Liggaamsmassa(kg)	Dosage/ Dosis	Grootte
Small	0,5 – 2	¼ tablet	Klein
	2 – 5	½ tablet	
	6 – 10	1 tablet	
Medium	11 – 15	1 ½ tablets/tablette	Middelslag
	16 – 20	2 tablets/tablette	
	21 – 25	2 ½ tablets/tablette	
	26 – 30	3 tablets/tablette	
Large	31 – 35	3 ½ tablets/tablette	Groot
	36 – 40	4 tablets/tablette	
Breed		Size/Grootte	Ras
Pekingese, Maltese, Pug, Chihuahua and puppies		Small/Klein (0,5 – 10 kg)	Pekinees, Maltees, Mopshond, Chihuahua en jong hondjies
Spaniel, Corgi and larger breed puppies		Medium/Middelslag (11 – 30 kg)	Patryshond, Korgi en groter ras jong hondjies
Rottweiler, Dobermann, German Shepherd and Great Dane		Large/Groot (31 – 40 kg)	Rottweiler, Dobermann, Duitse Herdershond en Deense hond
St. Bernard		Extra large/Extra groot (over 40 kg/meer as 40 kg)	St. Bernard
<ul style="list-style-type: none"> Puppies should be dewormed every 2 weeks, from the age of 2 – 12 weeks, and at regular intervals of 3 – 6 months thereafter to control worm infestation. Can be given safely to all dogs including old animals, young pups, working dogs, greyhounds and even to pregnant bitches 10 days before whelping. 		<ul style="list-style-type: none"> Ontworming van jong hondjies behoort elke 2 weke, van die ouderdom van 2 – 12 weke, plaas te vind en daarna met gereelde tussenposes van 3 – 6 maande om wurmbesmetting te beheer. Kan met veiligheid toegedien word aan ou diere, jong hondjies, werkende honde, windhonde asook dragtige tewe tot 10 dae voor welp. 	
Efficacy Controls: ≥ 90 % effective	Species/Spesie	Doeltreffendheid Beheer: ≥ 90 % doeltreffend	
Ascarid	<i>Toxocara canis</i>	Askariede	
Hookworm	<i>Ancylostoma caninum</i>	Haakwurm	
Tapeworms	<i>Dipylidium caninum</i> <i>Taenia</i> spp. <i>Echinococcus</i> spp.	Lintwurms	
Whipworm	<i>Trichuris vulpis</i>	Sambokwurm	

Registration holder/Registrasiehouer:

Ascendis Vet (Pty) Ltd
 Co. Reg. No. 2001/017471/07
 31 Georgian Crescent East
 Bryanston
 Johannesburg
 South Africa
 2191
 Tel: [+27 10 880 0778](tel:+27108800778)
www.ascendisanimalhealth.com